

Cusp Singularities in the Magnetization Curve of Frustrated Kondo Necklace Model

Takahiro Yamamoto *, Rumi Manago, Yoshihiro Mori and Chikara Ishii

Department of Physics, Faculty of Science, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

(Received February 1, 2008)

Magnetization processes of frustrated Kondo necklace model are studied by means of a density matrix renormalization group (DMRG) method and an elementary band theory based on a bond-operator formalism. The DMRG calculations clearly show the cusp singularity in a low-magnetization region ($0 < m < 1/2$) besides that in a high-magnetization region ($1/2 < m < 1$) which is expected from previous studies on the magnetization curve of the Majumdar-Ghosh model. An appearance mechanism of the low-magnetization cusp is interpreted in terms of a double-well shape of a low-energy band arising from frustrations between nearest- and next-nearest-neighbor interactions. We also discuss critical behaviors of magnetization near the cusp and obtain a phase diagram showing whether the cusp appears in the magnetization curve or not.

KEYWORDS: magnetization curve, cusp singularity, frustration, DMRG, bond-operator

1. Introduction

Over the last few years, a large number of materials exhibiting quantum phase transitions induced by an external magnetic field have been discovered, and their critical properties have been attracted much attention experimentally as well as theoretically. Such field-induced phase transitions are associated with various singularities in the magnetization curve, such as plateaus,¹⁻⁴ cusps⁵⁻¹¹ and jumps.¹²⁻¹⁴ In contrast to concern about plateaus and jumps, very few attempts have been made at the problem of cusp singularity in the magnetization curve. The cusp singularity was first found in the SU(3) integrable spin chain known as the Lai-Sutherland model by Parkinson with the use of the Bethe ansatz method.⁵ Subsequently, it has been investigated with a focus on integrable models. This may be because it is difficult to find cusp singularities of non-integrable models by conventional numerical approaches like an exact-diagonalization and quantum Monte Carlo methods.

Very recently, Okunish *et al.* showed that the cusp singularities appear in the magnetization curve of non-integrable spin systems, such as the spin-1/2 zig-zag ladder and the spin-1 bilinear-biquadratic chain, using the product-wave-function renormalization group (PWFRG) method,^{15,16} which is a variant of the density-matrix renormalization group (DMRG) method.¹⁷ They also pointed out that the appearance of cusps originate in the double-well shape of the energy dispersion of the quasi-particles such as magnon, spinon excitations.

In this paper, we introduce *frustrated Kondo necklace model*, which is equivalent to the spin-1/2 chain with nearest- and next-nearest-neighbor interactions (J_1 and J_2) coupled to a localized spin on each site, as a new non-integrable spin system exhibiting the cusp singularities in its magnetization curve. Here, let us briefly review the magnetization process of the frustrated Kondo necklace model already understood by previous studies.¹⁸⁻²⁰ The low-energy excitations of this system have a finite gap

(so-called Kondo singlet gap), reflecting that the ground state in the absence of the magnetic field is always in a quantum disordered phase whose spin configuration is an aggregate of intra-atomic Kondo singlets.²¹⁻²⁵ When a magnetic field, h , is applied to the system, the magnetization of this system begins to grow at the Kondo singlet gap. As for the Kondo necklace model without J_2 ,²⁶ the magnetization curve smoothly grows with increasing the magnetic field, h , once the Kondo singlet gap is collapsed.¹⁸ On the other hand, in the presence of the strong frustration ($J_2/J_1 > 0.3419$), the magnetization plateau appears at half saturation ($m = 1/2$) besides the Kondo singlet gap.^{19,20} The $m = 1/2$ plateau can be explained as follows. The frustrated Kondo necklace model with $m = 1/2$ can be effectively regarded as spin-1/2 zig-zag ladder which is well known as the Majumdar-Ghosh model²⁷ in a special case $J_2/J_1 = 0.5$, because the innercore spins of the system are almost ferromagnetically aligned by the magnetic field h and their degree of freedom is almost frozen by h . It is well known that the ground state of spin-1/2 zig-zag ladder is in the quantum disordered singlet-dimer phase. This phase corresponds to the $m = 1/2$ plateau phase in the present system.

Since the magnetization curve of the spin-1/2 zig-zag ladder exhibits the cusp singularity in the vicinity of the saturation magnetization, we can expect that the magnetization curve of the present model exhibits a similar cusp singularity in the high-field region $1/2 < m < 1$. The expected cusp will be found in the high-field region with the use of DMRG method in the next section. Moreover, the DMRG calculations will find an unexpected cusp just above the Kondo singlet gap. An appearance mechanism of the low-field cusp differs from that of the high-field cusp, because the excitations of innercore spins play an essential role of the magnetization process in this region. The main purpose of this paper is to explain the origin and characteristics of the low-field cusp singularity.

This paper is organized as follows. In Sec. 2, the Hamiltonian of the frustrated Kondo necklace model will be

*E-mail address: takahiro@rs.kagu.tus.ac.jp




Fig. 1. Schematic representation of the frustrated Kondo necklace model (1). Small and Large circles denote the conduction electron spin (S_c) and the innercore spin (S_f) respectively.

given and its magnetization curves will be calculated by the DMRG method. In Sec. 3, the frustrated Kondo necklace model will be mapped onto a fermionic quasiparticle system by means of the bond-operator representation which can explicitly describe the triplet excitation from the Kondo singlet sea (Sec. 3.1). A mechanism of appearance of the low-field cusp will be explained in terms of the double-well shape of an energy dispersion curve of the triplet excitation (Sec. 3.2). Moreover we will discuss the critical behavior of the magnetization curve near the cusp singularity in the low-field region $0 < m < 1/2$ (Sec. 3.3). In Sec. 4, we will determine the phase diagram on a plane of the Kondo interaction, J_K , and the frustration parameter, $\lambda \equiv J_2/J_1$, showing whether the cusp singularity appears in the magnetization curve or not. The final section (Sec. 5) is devoted to conclusion.

2. Hamiltonian and magnetization curves

The Kondo necklace model is simplified version of the Kondo lattice model in one dimension, assuming a narrow band limit for conduction electrons in the Kondo lattice model. In this limit, the motion of conduction electrons is prohibited, and the system is composed of two different kinds of localized spins: the spins of innercore electrons (on the f orbits) and the conduction electrons. Thus, the Hamiltonian of frustrated Kondo necklace is described as

$$\mathcal{H} = J \sum_{j=1}^L \sum_{\nu=1,2} \lambda^{\nu-1} \mathbf{S}_{c,j} \cdot \mathbf{S}_{c,j+\nu} + J_K \sum_{j=1}^L \mathbf{S}_{c,j} \cdot \mathbf{S}_{f,j}, \quad (1)$$

which is schematically shown in Fig. 1. Here, $\mathbf{S}_{c,j}$ are conduction-electron spins with $S_{c,j} = 1/2$, and $\mathbf{S}_{f,j}$ are innercore-electron spins with $S_{f,j} = 1/2$ which are hanging from the S_c -spin chain. The nearest-neighbor exchange coupling J and Kondo coupling J_K are assumed to be antiferromagnetic ($J, J_K > 0$). λ is a frustration parameter which denotes competition between nearest- and next-nearest-neighbor interactions in S_c -spins.

In the calculation of the magnetization, we use the DMRG method based on the finite-size algorithm.¹⁷⁾ The advantage of this method is that the large system can be studied, as compared with the exact-diagonalization method. Unlike the quantum Monte Carlo simulations, the DMRG method is free from the negative sign problem and can easily calculate even the ground state in the magnetic field. The DMRG method is an ideal tool for studying the magnetization curve of the one-dimensional spin systems. Our DMRG calculations have been per-




Fig. 2. The magnetization curves of the frustrated Kondo necklace model with $J/J_K = 1.0$. In the case of $\lambda = 0.5$ (filled circle), two cusps and a plateau appear in the magnetization curve which is smooth in the absence of frustration $\lambda = 0.0$ (open circle).

formed under the open boundary condition, for the system length $L = 60$ and the number of optimal states up to 120.

Figure 2 shows the magnetization curves of the frustrated Kondo necklace model with $\lambda = 0.0$ and 0.5 for fixed $J/J_K = 1.0$. In the absence of frustration ($\lambda = 0.0$), the magnetization curve grows smoothly with increasing the magnetic field h once the Kondo singlet gap Δ_K collapses. This means that there is no phase transition in the magnetic field in the range $\Delta_K < h < h_S$ (h_S : saturation field) and the system belongs to a universality class of the Tomonaga-Luttinger liquid phase.^{19,20)}

In the case of $\lambda = 0.5$, the Fig. 2 clearly shows the cusp singularities in the low-field ($0 < m < 1/2$) and high-field ($1/2 < m < 1$) regions, which cannot be detected by the exact-diagonalization calculations. As we mentioned in Sec. 1, the appearance of high-field cusp is expected from previous results by Okunishi *et al.*,¹⁵⁾ and can be interpreted in terms of the low-energy dispersion with double-well shape of single down-spin magnon in the up-spin sea. However, the concept of single down-spin magnon is useful only in the vicinity of the saturation magnetization. Therefore it cannot be used to explain an appearance mechanism of the low-field cusp. To discuss the low-field cusp, we must consider appropriate elementary excitations in the low-field region.

3. Mechanism of an appearance of a cusp singularity

3.1 Bond-operator representation

In this section, we discuss an origin of cusp singularity in the low-field region ($0 < m < 1/2$). First of all, it should be remarked that we have to know about the low-energy excitation modes of the system, in order to understand the magnetization process in the low-field region. As was stressed before, the ground state of the present system (1) in the absence of the magnetic field is always in the Kondo singlet phase, when the Kondo coupling, J_K , of any strength is present. This suggests that the

low-energy excitations will be well described by triplet excitations from the Kondo singlet sea. To express this picture in an explicit form, we introduce the following bond-operators creating singlet and triplet states out of the vacuum $|0\rangle$:

$$s^\dagger |0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle), \quad (2)$$

$$t_x^\dagger |0\rangle = -\frac{1}{\sqrt{2}} (|\uparrow\uparrow\rangle - |\downarrow\downarrow\rangle), \quad (3)$$

$$t_y^\dagger |0\rangle = \frac{i}{\sqrt{2}} (|\uparrow\uparrow\rangle + |\downarrow\downarrow\rangle), \quad (4)$$

$$t_z^\dagger |0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle). \quad (5)$$

These bond-operators were first proposed by Sachdev and Bhatt²⁸⁾ in order to treat the quantum phase transitions between dimerized and magnetically ordered phases of antiferromagnets. Here, $\uparrow, \downarrow$ and $\uparrow, \downarrow$ denote z -components of S_c and S_f spins, respectively. To describe the magnetization process of the Kondo necklace model, it is more convenient to use another set of bond-operators $u^\dagger$ and $d^\dagger$ creating the states $|\uparrow\uparrow\rangle$ and $|\downarrow\downarrow\rangle$:

$$u^\dagger |0\rangle \equiv \frac{-1}{\sqrt{2}} (t_x^\dagger + it_y^\dagger) |0\rangle = |\uparrow\uparrow\rangle, \quad (6)$$

$$d^\dagger |0\rangle \equiv \frac{1}{\sqrt{2}} (t_x^\dagger - it_y^\dagger) |0\rangle = |\downarrow\downarrow\rangle, \quad (7)$$

instead of the transverse triplet operators t_x and t_y .²⁹⁾

In terms of these bond-operator representation, we can imagine that each lattice point of the Kondo necklace model is in one of the four atomic states $s^\dagger |0\rangle$, $u^\dagger |0\rangle$, $t_z^\dagger |0\rangle$, and $d^\dagger |0\rangle$, or equivalently, each site is occupied by one of the virtual particles created by bond-operators $s^\dagger$, $u^\dagger$, $t_z^\dagger$ and $d^\dagger$. Then, these operators should satisfy the local constraint,

$$s_j^\dagger s_j + u_j^\dagger u_j + t_{z,j}^\dagger t_{z,j} + d_j^\dagger d_j = 1, \quad (8)$$

in order to work in the physical subspace with a single particle on each site. The spin operators $S_{p,j}$ ($p = c, f$) are represented in terms of these bond-operators as

$$S_{p,j}^+ = \frac{1}{\sqrt{2}} (\pm s_j^\dagger d_j \mp u_j^\dagger s_j + t_{z,j}^\dagger d_j + u_j^\dagger t_{z,j}), \quad (9)$$

$$S_{p,j}^- = \frac{1}{\sqrt{2}} (\pm d_j^\dagger s_j \mp s_j^\dagger u_j + d_j^\dagger t_{z,j} + t_{z,j}^\dagger u_j), \quad (10)$$

$$S_{p,j}^z = \frac{1}{2} (\pm s_j^\dagger t_{z,j} \pm t_{z,j}^\dagger s_j - d_j^\dagger d_j + u_j^\dagger u_j), \quad (11)$$

where the signs in the upper and lower rows denote the cases of $p = c$ and f , respectively.

Thus far, we have not specified the algebra of the bond-operators. As a matter of fact, the spin algebra:

$$[S_{p,i}^+, S_{q,j}^-] = 2S_j^{p,z} \delta_{i,j} \delta_{p,q}, \quad [S_{p,i}^\pm, S_{q,j}^\pm] = \mp S_j^{p,\pm} \delta_{i,j} \delta_{p,q}, \quad (12)$$

can be derived by using the expressions (9)-(11), irrespective that the bond-operators obey either bosonic or fermionic commutation relations. It is usual to assume for the bond-operators to be bosonic, in order to perform a physically meaningful mean-field approximation

on the basis of Bose-Einstein condensation of singlet states, $\langle s \rangle = \langle s^\dagger \rangle \neq 0$, $\langle t \rangle = \langle t^\dagger \rangle = 0$. The bond-operator mean-field theory was successfully applied to the study of disordered dimerized phases in various quantum spin systems (e.g. spin-1/2 ladder,³⁰⁾ Bilayer Heisenberg model³¹⁾ and standard Kondo necklace model²¹⁾). Recently, the method of bosonic bond-operators was applied to the studies of magnetization process of spin-1 antiferromagnetic Heisenberg chain and spin-1/2 and spin-1 Heisenberg ladders, mapping onto spin-1/2 pseudospin Hamiltonian in the effective magnetic field.²⁹⁾

In closing this section, we note that the algebra of the bond-operators is not important in the following discussion. An important point in this work is that the triplet excitations from the Kondo singlet sea are represented as *fermionic* particles, as shown in next section. Thus, we impose here the bosonic commutation relation on the bond-operators, only for convenience. Of course, even if we impose the fermionic commutation relation to the bond-operators, our results in this paper are not changed.

3.2 Effective band theory

We are now in the position to state a reasonable picture of the origin of the cusp singularity in the low-field region, $0 < m < 1/2$, on the basis of an elementary band theory constructed with the use of bond-operator representation. To do this, the triplet excitations from the Kondo singlet sea are treated as *fermionic* particles.

Since the ground state is always in the Kondo singlet phase as mentioned before, it seems to be useful to start from the strong Kondo-coupling limit, $J/J_K \ll 1$. In the limit of strong- J_K and $\hbar \approx \Delta_K$, we can project out the states $t_z^\dagger |0\rangle$ and $d^\dagger |0\rangle$, preserving only two low-lying states $s^\dagger |0\rangle$ and $u^\dagger |0\rangle$. Then, the constraint (8) can be rewritten as

$$s_j^\dagger s_j + u_j^\dagger u_j = 1, \quad (13)$$

and the spin operators in Eqs. (9)-(11) are simplified as

$$S_{p,j}^+ = \mp \frac{1}{\sqrt{2}} u_j^\dagger s_j \equiv \mp \frac{1}{\sqrt{2}} \gamma_j^\dagger, \quad (14)$$

$$S_{p,j}^- = \mp \frac{1}{\sqrt{2}} s_j^\dagger u_j \equiv \mp \frac{1}{\sqrt{2}} \gamma_j, \quad (15)$$

$$S_{p,j}^z = \frac{1}{2} u_j^\dagger u_j \equiv \frac{1}{2} \gamma_j^\dagger \gamma_j. \quad (16)$$

The new operator $\gamma_j^\dagger \equiv u_j^\dagger s_j$ creates a triplet particle on j -th site from the Kondo singlet sea, satisfying the following *hardcore bosonic* commutation relations

$$\{\gamma_j, \gamma_j^\dagger\} \equiv \gamma_j \gamma_j^\dagger + \gamma_j^\dagger \gamma_j = 1, \quad (17)$$

$$(\gamma_j^\dagger)^2 = \gamma_j^2 = 0, \quad (18)$$

$$[\gamma_i, \gamma_j^\dagger] = [\gamma_i^\dagger, \gamma_j] = [\gamma_i, \gamma_j] = 0, \quad i \neq j. \quad (19)$$

Note that the operators $\gamma_j^\dagger$ and γ_j are neither fermionic nor bosonic, and thanks to Eq. (17), the constraint (13) is satisfied automatically. The hardcore bosons can be mapped onto spinless fermions through the Jordan-

Wigner transformation,

$$\xi_j^\dagger = \gamma_j^\dagger \exp \left[i\pi \sum_{n=1}^{j-1} \gamma_n^\dagger \gamma_n \right], \quad (20)$$

$$\xi_j = \exp \left[-i\pi \sum_{n=1}^{j-1} \gamma_n^\dagger \gamma_n \right] \gamma_j. \quad (21)$$

It is convenient to introduce the Fourier transform of fermionic operator,

$$\xi_j = \frac{1}{\sqrt{L}} \sum_k \xi_k \exp(ikj), \quad (22)$$

where the wave number k takes discrete values $k = 2\pi n/L$ ($n = 0, 1, \dots, L-1$) assuming L to be even integer. Using the fermionic operators $\xi_k^\dagger$ and ξ_k , the frustrated Kondo necklace model [Eq. (1)] in the magnetic field h is mapped onto a *fermionic* quasiparticle system:

$$\begin{aligned} \mathcal{H} = & E_0 + \sum_k (\epsilon_k - h) \xi_k^\dagger \xi_k \\ & + \frac{J}{4} \sum_{k,k',q} \cos(k+k'+q) \xi_{k+q}^\dagger \xi_{k'-q}^\dagger \xi_k \xi_{k'} \end{aligned} \quad (23)$$

where $E_0 = -3J_K L/4$ is an energy of the Kondo singlet sea,

$$|\text{vac}\rangle = s_1^\dagger s_2^\dagger \cdots s_L^\dagger |0\rangle, \quad (24)$$

satisfying a relation $\xi_k |\text{vac}\rangle = 0$ and the single-particle energy dispersion is given as

$$\epsilon_k = J_K + \frac{J}{2} [\cos k + \lambda \cos(2k)]. \quad (25)$$

The last term in Eq. (23) describes the scattering between the quasiparticles. This is negligible in the low-density region $\sum_k \langle \xi_k^\dagger \xi_k \rangle \ll 1$, corresponding to the low-magnetization $m \ll 1$.

We discuss the single-particle energy dispersion (25) in some detail. The dashed curves in Fig. 3 show the single-particle energy dispersion ϵ_k/J_K for $\lambda = 0.2$ and 0.4 where $J/J_K = 0.1$ is assumed. As is seen in Fig. 3 and Eq. (25), the shape of dispersion curve changes at the critical frustration $\lambda_c = 1/4$. (Taking account of higher order corrections, this value of the critical point $\lambda_c = 1/4$ will be modified later.) For $\lambda < 1/4$, the dispersion has a single bottom at $k = \pi$. On the other hand, for $\lambda > 1/4$, the dispersion curve has double-well shape with two bottoms at the wave number $\pi \pm \sin^{-1}(-1/4\lambda)$ ($0 \leq k < 2\pi$) and at the energy $\epsilon_b/J_K = 1 - (J/2J_K)(\lambda + 1/8\lambda)$. The state at $k = \pi$, being global minimum for $\lambda < 1/4$, turns out to be local maximum with the energy $\epsilon_m/J_K = 1 - (J/2J_K)(\lambda - 1)$.

These results can be also obtained by the perturbation expansion up to the first order in J/J_K , and provide a sufficient basis to interpret the origin of the appearance of the cusp qualitatively. More detailed analysis up to the second order in J/J_K is described in the Appendix, and the results are shown by solid curves in Fig. 3. As is seen from these plots, the second order results are in excellent agreement with those of exact diagonalization indicated by open circles.




Fig. 3. The energy dispersion of a single triplet fermion for $J/J_K = 0.1$ with $\lambda = 0.2$ (above) and 0.4 (below). Short-dash curve shows the dispersion given by Eq. (25). Solid curve represent the dispersion obtained by the perturbation expansion up to second order in J/J_K (see Appendix). The open circles show the exact-diagonalization data at $k = 2\pi n/L$ ($n = 0, 1, \dots, L-1$) for $L = 8, 10$ and 12 .

The eigenstate with the energy ϵ_k is given as a single quasiparticle state,

$$\begin{aligned} |k\rangle &= \xi_k^\dagger |\text{vac}\rangle \\ &= \frac{1}{\sqrt{L}} \sum_{j=1}^L \exp(ikj) s_1^\dagger s_2^\dagger \cdots u_j^\dagger \cdots s_L^\dagger |0\rangle. \end{aligned} \quad (26)$$

To study the low-energy properties, one often makes the assumption that additional many quasiparticle eigenstates can be constructed as a collection of independent quasiparticles, that is $|k_1, k_2, \dots, k_N\rangle = \xi_{k_1}^\dagger \xi_{k_2}^\dagger \cdots \xi_{k_N}^\dagger |0\rangle$ with the energy $\epsilon_{k_1} + \epsilon_{k_2} + \cdots \epsilon_{k_N}$.

We have prepared now to interpret the low-field magnetization process just above the Kondo singlet gap Δ_K in terms of elementary band theory with the use of above single-particle energy dispersion relation (25). Thereby, the magnetic field h plays a role of the chemical potential or the Fermi energy and the magnetization m corresponds to the quasiparticle number per site occupying the state $|k\rangle$ below the Fermi energy h . Thus the magnetization at zero temperature is given as

$$m = \frac{1}{L} \sum_{|k| < k_F} \langle \xi_k^\dagger \xi_k \rangle \rightarrow \frac{1}{2\pi} \int_0^{2\pi} \theta(h - \epsilon_k) dk, \quad (27)$$

where k_F and $\theta(x)$ are the Fermi wave-number and the




Fig. 4. Schematic diagram showing the relation between the double-well shaped low-energy dispersion of triplet quasiparticle and the cusp singularity in the low-field magnetization curve. ‘G.S.’ denotes the energy of ground state.

Heaviside’s step function, respectively.

On the basis of Eq. (27), we can explain the characteristic features of magnetization curve in the low-field region, shown in Fig. 4. When a magnetic field h is applied to the system, no magnetization appears until h arrives at the bottoms of the dispersion. If the frustration parameter λ is larger than the critical value λ_c in the region $\epsilon_m > h > \epsilon_b$, the Fermi level crosses the dispersion at four wave-numbers, satisfying $\epsilon_k = h$, and the magnetization curve shows sharp rise. With further increase in the magnetic field, the Fermi level eventually arrives at the local maximum ϵ_m at $k = \pi$, and the number of crossing points of the Fermi level and the dispersion curve suddenly reduces from 4 to 2, resulting in a sudden slowdown of the rate of band filling and growth rate of the magnetization. This sudden change in the rate of band filling is reflected as the cusp singularity on the magnetization curve of the Kondo necklace model.

3.3 Critical behavior near the cusp singularity

To discuss the detailed behavior of the magnetization curve in the vicinity of cusp singularity, we consider the field derivative of the magnetization dm/dh . From Eq. (27), we find that dm/dh is given by the quasiparticle density of states at the Fermi energy h as follows.

$$\begin{aligned} \frac{dm}{dh} &= \frac{1}{2\pi} \int_0^{2\pi} \delta(h - \epsilon_k) dk \\ &= \frac{1}{2\pi} \sum_{\{\epsilon_k=h\}} \left| \frac{d\epsilon_k}{dk} \right|^{-1}. \end{aligned} \quad (28)$$

Here, $\sum_{\{\epsilon_k=h\}}$ denotes a sum over all Fermi points. When the energy dispersion is shaped like the double-well, two different situations may occur (see Fig. 5(a)).

[1] For $h > \epsilon_m$, there are two Fermi points $\pm k_F$, and the energy dispersion can be linearized around the Fermi points as far as low-energy excitations are concerned: $\epsilon_k = \pm v_F \{(k - \pi) \mp k_F\} + h$, where v_F is the Fermi velocity of the ξ_k -fermions. Thus, from Eq. (28), we obtain $dm/dh = L/\pi v_F$ which is almost independent of h




Fig. 5. (a) The effective low-energy dispersion. The dashed lines show the magnetic field h corresponding to the Fermi level. (b) The field derivative of the magnetization dm/dh near the critical point $h = \epsilon_m$ of the frustrated Kondo necklace model with $J/J_K = 1.0$ and $\lambda = 0.5$. The solid line denotes asymptotic curve $dm/dh \sim (\epsilon_m - h)^{-1/2}$ near $h = \epsilon_m$.

because v_F only depends weakly on h .

[2] For $h < \epsilon_m$, there are four Fermi points satisfying the relation, $2k_{F1} - 2k_{F2} = m$. In this case, low-energy states are effectively described by the two-band model [See Fig. 5(a)]:

$$\epsilon_k^{(1)} = \pm v_{F1} \{(k - \pi) \mp k_{F1}\} - h, \quad (29)$$

$$\epsilon_k^{(2)} = -C(k - \pi)^2 + \epsilon_m, \quad (30)$$

where C is a positive constant. The parabolic dependence of $\epsilon_k^{(2)}$ around $k = \pi$ gives decisive difference in dm/dh from the case of [1]. Substituting Eqs. (29) and (30) for Eq. (28), we obtain the most dominant contribution

$$\frac{dm}{dh} \sim \frac{L}{2\pi\sqrt{C}} \frac{1}{\sqrt{\epsilon_m - h}} \quad (31)$$

which diverges at $h = \epsilon_m$. Thus, we find that the critical exponent characterizing the intensity of the divergence of the field derivative of the magnetization, dm/dh , is given as $1/2$.

Plots in Fig. 5(b) show dm/dh of the frustrated Kondo necklace model (1) with $J/J_K = 1.0$ and $\lambda = 0.5$, obtained by numerically differentiating the DMRG data in Fig. 2. As is expected from the above results, for $h > \epsilon_m$, the plots depend on h very weakly. On the other hand, for $h < \epsilon_m$, it diverges as $dm/dh \sim (\epsilon_m - h)^{-1/2}$ as $h \rightarrow \epsilon_m$ (the solid curve in Fig. 5(b)). Away from the cusp singular point $h = \epsilon_m$, the data are almost independent of h . This means that the low-energy dispersion is described by two linear-bands involving four Fermi points. From the above results, we find that the magnetization behaves as

$$m - m_{\text{cusp}} \propto \begin{cases} h - \epsilon_m & : h > \epsilon_m \\ -\sqrt{\epsilon_m - h} & : h < \epsilon_m \end{cases} \quad (32)$$

in the vicinity of the critical point $h = \epsilon_m$ with $m = m_{\text{cusp}}$. This result well explains the behavior of magnetization curves near the cusp singularity obtained by the DMRG calculations (See Fig. 2).

4. Phase Diagram

In the previous sections, we saw that the appearance of low-field cusp singularity in the magnetization curve is governed by the competition between the Kondo coupling J/J_K and frustration parameter λ . In the strong Kondo coupling limit $J/J_K \rightarrow 0$, the cusp singularity appears when λ exceeds the value $\lambda_c = 1/4$. In the presence of Kondo coupling of general strength, the value of λ_c will depend on the strength of Kondo coupling. In the following, we determine a phase boundary on the plane of J/J_K and λ , dividing whether the cusp appears in the magnetization curve or not. To do this, we note that the value of energy dispersion curve at zone-center point $k = \pi$ turns from global minimum to local maximum. Therefore, the critical point λ_c is to be determined from the condition that the point $k = \pi$ is the deflection point of the energy dispersion:

$$\left(\frac{\partial^2 \epsilon_k}{\partial k^2} \right)_{k=\pi} = 0. \quad (33)$$

As shown in Appendix, the critical value of frustration λ_c , including the second order correction with respect to J/J_K is given as

$$\lambda_c = \frac{2J_K + 2J}{8J_K + 5J} \approx \frac{1}{4} + \frac{3}{32} \left(\frac{J}{J_K} \right) - \frac{15}{256} \left(\frac{J}{J_K} \right)^2. \quad (34)$$

In order to determine λ_c outside the strong Kondo-coupling region, however, we need the expression of energy dispersion including the full effects of Kondo cou-

pling and frustration. This is performed by determining the energy dispersion for various values of J/J_K by means of the exact-diagonalization method.

Here we briefly explain our exact-diagonalization method. Our exact-diagonalization calculations were performed under the periodic boundary condition $S_{p,j} = S_{p,j+L}$ ($p = c$ and f). Since the z -component of total spin operator commutes with the frustrated Kondo necklace Hamiltonian (1), all eigenvalues and eigenstates of the Hamiltonian are labeled by the total magnetization $M = \sum_{j=1}^L (\sigma_j^z + S_j^z)$. In addition, since the Hamiltonian (1) is invariant under the translation along the chain-direction ($S_{p,j} \rightarrow S_{p,j+1}$), the eigenvalues and eigenstates are classified by the wave number $k = 2\pi n/L$ ($n = 0, 1, \dots, L-1$) corresponding to eigenvalue of the translation operator. Thus, in Fig. 3, we plot the energy dispersion, $E(k, M=1) - E(k, M=0)$ with $k = 2\pi n/L$, for three lengths $L = 8, 10$, and 12 .

Since the energy dispersion is an even function around $k = \pi$, we fit the exact-diagonalization data near $k = \pi$ using a fitting function,

$$\epsilon_k = \Delta_K + \frac{\alpha}{2!}(k - \pi)^2 + \frac{\beta}{4!}(k - \pi)^4 + \dots, \quad (35)$$

up to eighth order and estimate fitting parameters Δ_K , α , β , $\dots$ for various λ and fixed J/J_K . The parameter $\alpha = (\partial^2 \omega / \partial k^2)_{k=\pi}$ thus obtained for $J/J_K = 0.1, 0.3$ and 0.5 are shown by circles, squares and triangles in Fig. 6, respectively. The next step is to interpolate α in order to obtain the critical point λ_c satisfying the condition (33). We assumed a liner form of interpolation function $\alpha = A(\lambda - \lambda_c)$ in analogy with that in the strong Kondo-coupling limit (see Appendix). Thus we can obtain the critical point λ_c from the condition $\alpha = 0$ (see Fig. 6). Finally, the phase diagram on $(J/J_K, \lambda)$ -plane is constructed, performing above series of procedures for various values of J/J_K , as shown in Fig. 7. The short-dash curve in Fig. 7 shows the boundary given by Eq. (34) and is in good agreement with the exact-diagonalization data in the strong Kondo-coupling region, $J/J_K \ll 1$. On the boundary, the magnetization curve grows rapidly at $h = \Delta_K$ because the low-energy part of the dispersion becomes quartic. To be more explicit, the magnetization behaves as $m \sim (h - \Delta_K)^{3/4}$, as is derived from Eq. (28).

5. Conclusion

In this paper, the magnetization processes of frustrated Kondo necklace model have been studied by using the density matrix renormalization group (DMRG) method. We have paid attention to the low-field cusp singularity appearing just above the Kondo singlet gap. To reveal an appearance mechanism of the cusp, we introduced the triplet quasiparticle excited from Kondo singlet sea by using the bond-operator representation of dimerized spins, and we have discussed the mechanism in terms of an elementary band theory with the use of a single-particle energy dispersion of the triplet quasiparticle. As a result, we have made it clear that the appearance of low-field cusp has its origin in a change of shape of quasiparticle energy dispersion curve, reflecting effects




Fig. 6. The second derivative of the energy dispersion ϵ_k at $k = \pi$. The circles, squares and triangles denote $(\partial^2 \epsilon_k / \partial k^2)_{k=\pi}$ for $J/J_K = 0.1, 0.3$ and 0.5 , respectively. Each symbol is fitted by a straight line.




Fig. 7. The phase diagram on plane of J/J_K and λ , which decides whether a cusp appears in the region of $0 < m < 1/2$ in the magnetization curve. The short-dash curve is the boundary given by Eq. (34) obtained from the strong Kondo-coupling expansion up to second order.

of frustration.

We also discussed the critical behavior of the magnetization in the vicinity of the cusp singularity in the low-field region. Calculating the density of states from the quasiparticle energy dispersion, we found that the magnetization curve behaves as $m - m_{\text{cusp}} \propto \sqrt{\epsilon_m - h}$ ($h < \epsilon_m$) and $m - m_{\text{cusp}} \propto h - \epsilon_m$ ($h > \epsilon_m$). This result well explains the behavior of the magnetization curves near the cusp obtained by the DMRG calculations. Moreover, we obtained the phase diagram on the $(J/J_K, \lambda)$ -plane deciding whether the cusp appears in the magnetization curve or not.

The effective band theory based on the bond-operator formalism must be successfully applied to various quantum spin systems with dimer-gapped ground state.³²⁾

Acknowledgments

We are grateful to M. Asano for his assistance in the DMRG calculations. We also would like to thank Profs. M. Suzuki, K. Hida, T. Nikuni, M. Nakamura and K. Okunishi for valuable comments and useful discussions. This work has been supported in part by Frontier Research Center for Computational Science (FRCCS) of Tokyo University of Science.

Appendix: Strong Kondo coupling expansion up to second order in J/J_K

In this appendix, we present the second order correction in J/J_K to the low-energy dispersion of the frustrated Kondo necklace model.

The Kondo coupling Hamiltonian [the second term in Eq. (1)] will be treated as the unperturbed one and its ground state is given as a direct product of the Kondo singlets [Eq. (24)]. On the other hand, intra-chain Hamiltonian [the first term in Eq. (1)] will be treated as the perturbation. The first order perturbation energy vanishes owing to the orthogonal condition. The second-order correction to the energy of ground state is calculated by the standard perturbation formula. The perturbation energy up to the second order in J/J_K is obtained as

$$E_0/J_K = -\frac{3L}{4} \left\{ 1 + \frac{1}{8}(1 + \lambda^2) \left(\frac{J}{J_K} \right)^2 \right\}. \quad (\text{A}\cdot 1)$$

To obtain the low-energy dispersion, we next calculate the energy correction in an excited state. We suppose that the low-energy physics in the strong Kondo-coupling region is well described by the single triplet excitation state (26). Following the same procedure as calculating the ground-state energy, we can estimate the excitation energy. The first and the second order energies are given as

$$\begin{aligned} E_k/J_K &= \left(-\frac{3}{4}L + 1 \right) + \frac{J}{2J_K} (\cos k + \lambda \cos(2k)) \\ &\quad - \frac{1}{32} \left(\frac{J}{J_K} \right)^2 \{ (1 + \lambda^2)(3L + 2) \\ &\quad - 4(2 - \lambda) \cos(k) + 2(1 - 4\lambda^2) \cos(2k) \\ &\quad + 4\lambda \cos(3k) + 2\lambda^2 \cos(4k) \}. \end{aligned} \quad (\text{A}\cdot 2)$$

Subtracting Eq. (A·2) from Eq. (A·1), we obtain the single-particle energy dispersion up to the second order in J/J_K :

$$\begin{aligned} \epsilon_k/J_K &\equiv (E_k - E_0)/J_K \\ &= 1 - \frac{1}{16} \left(\frac{J}{J_K} \right)^2 (1 + \lambda^2) \\ &\quad + \frac{J}{8J_K} \left\{ 4 + (2 - \lambda) \frac{J}{J_K} \right\} \cos k \\ &\quad + \frac{J}{16J_K} \{ 8\lambda - (1 - 4\lambda^2) \} \frac{J}{J_K} \cos(2k) \\ &\quad - \frac{\lambda}{8} \left(\frac{J}{J_K} \right)^2 \cos(3k) - \left(\frac{\lambda J}{4J_K} \right)^2 \cos(4k) \end{aligned}$$

where the first order correction is consistent with Eq. (25). The coefficients Δ_K , α and β of the Taylor expansion [Eq. (35)] of the energy dispersion (A.3) are given by

$$\Delta_K = 1 - \frac{J}{8J_K} (1 - \lambda) \left[4 + (3 + \lambda) \frac{J}{J_K} \right], \quad (\text{A.4})$$

$$\alpha = -\frac{J}{J_K} \left(2 + \frac{5J}{4J_K} \right) \left(\lambda - \frac{2J_K + 2J}{8J_K + 5J} \right), \quad (\text{A.5})$$

$$\beta = \frac{J}{J_K} \left[8\lambda - \frac{1}{2} - \frac{J}{4J_K} (48\lambda^2 - 41\lambda + 5) \right] \quad (\text{A.6})$$

Using the expression for α , we can determine an asymptotic curve of boundary on the plane of J/J_K and λ , dividing whether the cusp appears in the magnetization curve or not. Since the boundary, λ_c , is determined by the condition, $\alpha = 0$, we can obtain the boundary form (34) as shown in Fig. 7.

- 1) K. Hida, J. Phys. Soc. Jpn. **63** (1994) 2359.
- 2) K. Totsuka, Phys. Lett. A **228** (1997) 103; Phys. Rev. B **57** (1998) 3454.
- 3) M. Oshikawa, M. Yamanaka and I. Affleck, Phys. Rev. Lett. **78** (1997) 1984.
- 4) W. Shiramura, K. Takatsu, B. Kurniawan, H. Tanaka, H. Uekusa, Y. Ohashi, K. Takizawa, H. Mitamura and T. Goto J. Phys. Soc. Jpn. **67** (1998) 1548.
- 5) J. B. Parkinson, J. of Phys.: Cond. Matt. **1** (1989) 6709.
- 6) H. Kiwata, J. of Phys.: Cond. Matt. **7** (1995) 7991.
- 7) A. Schmitt, K.-H. Mütter and M. Karbach, J. of Phys. A **29** (1996) 3951.
- 8) H. Kiwata and Y. Akutsu, J. Phys. Soc. Jpn. **63** (1994) 4269.
- 9) P. Schlottmann, Phys. Rev. B **49** (1994) 9202.

- 10) A. A. Zvyagin and P. Schlottmann, Phys. Rev. B **52** (1995) 6569.
- 11) M. Fujii, S. Fujimoto and N. Kawakami, J. Phys. Soc. Jpn. **65** (1996) 2381.
- 12) M. Kohno and M. Takahashi, Phys. Rev. B **56** (1997) 3212.
- 13) C. Gerhardt, K.-H. Mütter and H. Kröger, Phys. Rev. B **57** (1998) 11504.
- 14) S. Hirata, cond-mat/9912066.
- 15) K. Okunishi, Y. Hieda and Y. Akutsu, Phys. Rev. B **60** (1999) R6953.
- 16) K. Okunishi and N. Maeshima, Phys. Rev. B **64** (2001) 212406.
- 17) S. R. White, Phys. Rev. Lett. **69** (1992) 2863; Phys. Rev. B **48** (1993) 10345.
- 18) T. Yamamoto, M. Asano and C. Ishii, J. Phys. Soc. Jpn. **70** (2001) 3678.
- 19) T. Yamamoto and C. Ishii, Prog. of Theor. Phys. Suppl. **145** (2002) 64.
- 20) T. Yamamoto, K. Ide and C. Ishii, Phys. Rev. B **66** (2002) 104408.
- 21) G.-M. Zhang, Q. Gu and L. Yu, Phys. Rev. B **62** (2000) 69.
- 22) R. T. Scalettar, D. J. Scalapino and R. L. Sugar, Phys. Rev. B **31** (1985) 7316.
- 23) S. Moukouri, L. G. Caron, C. Bourbonnais and L. Hubert, Phys. Rev. B **51** (1994) 15920.
- 24) H. Otsuka and T. Nishino, Phys. Rev. B **52** (1995) 15066.
- 25) Y. Chen, Q. Yuan, H. Chen and Y. Zhang, Phys. Lett. A **245** (1998) 167.
- 26) S. Doniach, Physica B **91** (1977) 231.
- 27) C. K. Majumdar and D. K. Ghosh, J. Math. Phys. **10** (1969) 1399.
- 28) S. Sachdev and R. N. Bhatt, Phys. Rev. B **41** (1990) 9323.
- 29) H.-T. Wang, H. Q. Lin and J.L. Shen, Phys. Rev. B **61** (2000) 4019.
- 30) S. Gopalan, T. M. Rice and M. Sigrist, Phys. Rev. B **49** (1993) 8901.
- 31) Y. Matsushita, M. P. Gelfand and C. Ishii, J. Phys. Soc. Jpn. **68** (1999) 247.
- 32) T. Yamamoto and C. Ishii, Physica B **329** (2003) 1277.